

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

Scientific name: Iris pseudacorus USDA Plants Code: IRPS
 Common names: Yellow iris
 Native distribution: Europe, Asia, northern Africa, and Mediterranean region
 Date assessed: 17 March 2008, 18 June 2008, 1 July 2008
 Assessors: Jinshuang Ma, Steven Clemants, Gerry Moore, Marilyn Jordan
 Reviewers: LIISMA SRC
 Date Approved: 9 July 2008; edited Dec. 15, 2008 Form version date: 22 October 2008

New York Invasiveness Rank: High (Relative Maximum Score 70.00-80.00)

| Distribution and Invasiveness Rank (<i>Obtain from PRISM invasiveness ranking form</i>) | | |
|--|----------------------|-------------------------|
| Status of this species in each PRISM: | Current Distribution | PRISM Invasiveness Rank |
| 1 Adirondack Park Invasive Program | Not Assessed | Not Assessed |
| 2 Capital/Mohawk | Not Assessed | Not Assessed |
| 3 Catskill Regional Invasive Species Partnership | Not Assessed | Not Assessed |
| 4 Finger Lakes | Not Assessed | Not Assessed |
| 5 Long Island Invasive Species Management Area | Common | High |
| 6 Lower Hudson | Not Assessed | Not Assessed |
| 7 Saint Lawrence/Eastern Lake Ontario | Not Assessed | Not Assessed |
| 8 Western New York | Not Assessed | Not Assessed |

| Invasiveness Ranking Summary (see details under appropriate sub-section) | | Total (Total Answered*) Possible | Total |
|--|---|---|-----------------|
| 1 | Ecological impact | 40 (<u>40</u>) | 24 |
| 2 | Biological characteristic and dispersal ability | 25 (<u>25</u>) | 19 |
| 3 | Ecological amplitude and distribution | 25 (<u>25</u>) | 21 |
| 4 | Difficulty of control | 10 (<u>10</u>) | 7 |
| | Outcome score | 100 (<u>100</u>) ^b | 76 ^a |
| | Relative maximum score † | | 76.00 |
| | New York Invasiveness Rank § | High (Relative Maximum Score 70.00-80.00) | |

* For questions answered "unknown" do not include point value in "Total Answered Points Possible." If "Total Answered Points Possible" is less than 70.00 points, then the overall invasive rank should be listed as "Unknown."

† Calculated as 100(a/b) to two decimal places.

§ Very High >80.00; High 70.00-80.00; Moderate 50.00-69.99; Low 40.00-49.99; Insignificant <40.00

A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

| | | |
|--|--|--|
| A1.1. Has this species been documented to persist without cultivation in NY? (reliable source; voucher not required) | | |
| <input checked="" type="checkbox"/> | Yes – continue to A1.2 | |
| <input type="checkbox"/> | No – continue to A2.1 | |
| A1.2. In which PRISMs is it known (see inset map)? | | |
| <input checked="" type="checkbox"/> | Adirondack Park Invasive Program | |
| <input checked="" type="checkbox"/> | Capital/Mohawk | |
| <input checked="" type="checkbox"/> | Catskill Regional Invasive Species Partnership | |
| <input checked="" type="checkbox"/> | Finger Lakes | |
| <input checked="" type="checkbox"/> | Long Island Invasive Species Management Area | |
| <input checked="" type="checkbox"/> | Lower Hudson | |
| <input checked="" type="checkbox"/> | Saint Lawrence/Eastern Lake Ontario | |
| <input checked="" type="checkbox"/> | Western New York | |

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

Documentation:

Sources of information:

Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

A2.1. What is the likelihood that this species will occur and persist outside of cultivation, given the climate in the following PRISMs? (obtain from PRISM invasiveness ranking form)

| | |
|--------------|--|
| Not Assessed | Adirondack Park Invasive Program |
| Not Assessed | Capital/Mohawk |
| Not Assessed | Catskill Regional Invasive Species Partnership |
| Not Assessed | Finger Lakes |
| Very Likely | Long Island Invasive Species Management Area |
| Not Assessed | Lower Hudson |
| Not Assessed | Saint Lawrence/Eastern Lake Ontario |
| Not Assessed | Western New York |

Documentation:

Sources of information (e.g.: distribution models, literature, expert opinions):

Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

If the species does not occur and is not likely to occur with any of the PRISMs, then stop here as there is no need to assess the species.

A2.2. What is the current distribution of the species in each PRISM? (obtain rank from PRISM invasiveness ranking forms)

| | |
|--|--------------|
| | Distribution |
| Adirondack Park Invasive Program | Not Assessed |
| Capital/Mohawk | Not Assessed |
| Catskill Regional Invasive Species Partnership | Not Assessed |
| Finger Lakes | Not Assessed |
| Long Island Invasive Species Management Area | Common |
| Lower Hudson | Not Assessed |
| Saint Lawrence/Eastern Lake Ontario | Not Assessed |
| Western New York | Not Assessed |

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2008; Weldy & Werier, 2005.

A2.3. Describe the potential or known suitable habitats within New York. Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

| | | |
|--|--|---|
| <p>Aquatic Habitats</p> <input type="checkbox"/> Salt/brackish waters <input checked="" type="checkbox"/> Freshwater tidal <input checked="" type="checkbox"/> Rivers/streams <input checked="" type="checkbox"/> Natural lakes and ponds <input type="checkbox"/> Vernal pools <input checked="" type="checkbox"/> Reservoirs/impoundments* | <p>Wetland Habitats</p> <input type="checkbox"/> Salt/brackish marshes <input checked="" type="checkbox"/> Freshwater marshes <input type="checkbox"/> Peatlands <input checked="" type="checkbox"/> Shrub swamps <input checked="" type="checkbox"/> Forested wetlands/riparian <input checked="" type="checkbox"/> Ditches* <input type="checkbox"/> Beaches and/or coastal dunes | <p>Upland Habitats</p> <input type="checkbox"/> Cultivated* <input type="checkbox"/> Grasslands/old fields <input type="checkbox"/> Shrublands <input type="checkbox"/> Forests/woodlands <input type="checkbox"/> Alpine <input type="checkbox"/> Roadsides* |
|--|--|---|

Other potential or known suitable habitats within New York:

Tu 2003 reports species in brackish habitats but we have no other evidence of it being salt tolerant. Moore has seen it in freshwater tidal zones but not in saline habitats.

Documentation:

Sources of information:

Goldblatt, 2002; Tu, 2003; Brooklyn Botanic Garden, 2008.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

B. INVASIVENESS RANKING

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score 7

Documentation:

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)

Can form a rhizome mat that collects sediment, elevates topography, and interferes with stream flow, resulting in a drier habitat.

Sources of information:

Thomas, 1980; Tu, 2003; Gravuer, 2005; Heckman, 2005.

1.2. Impact on Natural Community Structure

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score 3

Documentation:

Identify type of impact or alteration:

Can dominate the herb layer thus changing its density.

Sources of information:

Thomas, 1980; Tu, 2003; Gravuer, 2005; Heckman, 2005; S.Flint 2008 pers. comm.

1.3. Impact on Natural Community Composition

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) 10
- U. Unknown

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Score

| |
|---|
| 7 |
|---|

Documentation:

Identify type of impact or alteration:

The rhizome mats have been shown to alter microhabitats and prevent germination of native species (Hickman, 2005) and alter successional trajectories (Tu, 2003). Along the lower Potomac River, yellow iris's elevation of the seed bank is believed to have facilitated the conversion of riparian marshes into mesic forest (Thomas 1980). Greatly reduces abundance of native species and may ultimately eliminate natives in older infestations in the Adirondacks (S.FlintP. If similar impacts were documented from our area that would warrant the answer of D.

Sources of information:

Thomas, 1980; Tu, 2003; Heckman, 2005; S. Flint 2008 pers comm .

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades.

Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- | | | |
|----|--|----|
| A. | Negligible perceived impact | 0 |
| B. | Minor impact | 3 |
| C. | Moderate impact | 7 |
| D. | Severe impact on other species or species groups | 10 |
| U. | Unknown | |

Score

| |
|---|
| 7 |
|---|

Documentation:

Identify type of impact or alteration:

In Connecticut, *I. pseudacorus* has been shown to exclude the native arrow-arum (*Peltandra virginica*), a plant whose fruits are an important food for wildlife (Cox 1999, cited in Tu, 2003). It has also been shown to outcompete cat-tail (*Typha latifolia*) (Washington State Noxious Weed Control Board, 2003) and impact willows (*Salix* spp.) (Thomas, 1980). The species does not provide food for native wildlife species (Mehrhoff, 2003; Tu, 2003). However, sites in NY are not generally large.

Sources of information:

Thomas, 1980; Cox, 1999; Mehrhoff et al., 2003; Tu, 2003; Washington State Noxious Weed Control Board, 2003.

| | | | |
|----|-------------------|---|----|
| | Total Possible | <table border="1"><tr><td style="width: 40px; text-align: center;">40</td></tr></table> | 40 |
| 40 | | | |
| | Section One Total | <table border="1"><tr><td style="width: 40px; text-align: center;">24</td></tr></table> | 24 |
| 24 | | | |

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction (provisional thresholds, more investigation needed)

- | | | |
|----|---|---|
| A. | No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). | 0 |
| B. | Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) | 1 |
| C. | Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) | 2 |
| D. | Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not | 4 |

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

known, then maximum seed production reported to be greater than 1000 seeds per plant.)

U. Unknown

Score 4

Documentation:

Describe key reproductive characteristics (including seeds per plant):

Abundant reproduction and spread through abundant production of seeds and rhizome fragments, both of which are buoyant.

Sources of information:

Cody, 1961; Dalton, 1983; Ramsey and Peichel, 2001; Goldblatt, 2002; Tu, 2003; Washington State Noxious Weed Control Board, 2003; Weber, 2003; Gravuer, 2005; Heckman, 2005; ISSG, 2005.

2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score 4

Documentation:

Identify dispersal mechanisms:

Buoyant seeds and rhizomes could travel long distances, certainly beyond 100 meters.

Sources of information:

Tu, 2003; Gravuer, 2005; Heckman, 2005; ISSG, 2005.

2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

Score 3

Documentation:

Identify dispersal mechanisms:

Can be found for sale in nurseries, although it is not a popular horticultural plant since its flowering period is short. Boating activities might spread the seeds and rhizomes.

Sources of information:

Ramey & Peichel, 2001; Tu, 2003; Gravuer, 2005; Heckman, 2005; ISSG, 2005.

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- B. Possesses one characteristic that increases competitive advantage 3
- C. Possesses two or more characteristics that increase competitive advantage 6
- U. Unknown

Score 6

Documentation:
 Evidence of competitive ability:
 Shade tolerant, perennial, fast growth, fire tolerant.
 Sources of information:
 Ramey & Peichel, 2001; Tu, 2003; Washington State Noxious Weed Control Board, 2003; Weber, 2003; Gravuer, 2005; Heckman, 2005; ISSG, 2005.

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
- B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
- U. Unknown

Score 0

Documentation:
 Describe growth form:
 Can form large stands but does not exhibit a climbing or smothering habit, nor does it form dense thickets or mats.
 Sources of information:
 Tu, 2003; Washington State Noxious Weed Control Board, 2003; Gravuer, 2005; Heckman, 2005; ISSG, 2005; author's (Moore's) personal observations.

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
- B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
- C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
- U. Unknown (No studies have been completed)

Score 2

Documentation:
 Describe germination requirements:
 Germination of seeds in the field is rather low. Requires moist, but not inundated soil for germination but even here germination rates in the field were less than 20%. Scarification (such as heat from a fire) may increase germination rate. Seeds do appear to be tolerant of saline exposure.
 Sources of information:
 Dymes, 1920; Jesson, 1955; Thomas, 1980; Sutherland, 1990; Coops & Van Der Velde, 1995.

2.7. Other species in the genus invasive in New York or elsewhere

- A. No 0
- B. Yes 3
- U. Unknown

Score 0

Documentation:
 Species:
 Iris x conglomerata, Iris cristata, Iris verna var. smalliana, and Iris sibirica all reported from New York but none currently reported as invasive in NY or elsewhere.

Total Possible 25

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Section Two Total 19

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: “The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude”)

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score 4

Documentation:

Identify reason for selection, or evidence of weedy history:
 Large dense stands are usually present in disturbed areas with other invasive species in the NY metro area. However, in the Adirondacks yellow iris has rapidly invaded pristine areas with few/no other invasive species, and formed large stands (Bouquet River and Mt. Arab Lake; S. Flint 2008 pers comm).
 Sources of information:
 Cody, 1961; Dalton, 1983; Tu, 2003; Invasive Species Specialist Group 2005; author's (Moore's) personal observations; Steve Glenn's personal observations.

3.2. Number of habitats the species may invade

- A. Not known to invade any natural habitats given at A2.3 0
- B. Known to occur in two or more of the habitats given at A2.3, with at least one a natural habitat. 1
- C. Known to occur in three or more of the habitats given at A2.3, with at least two a natural habitat. 2
- D. Known to occur in four or more of the habitats given at A2.3, with at least three a natural habitat. 4
- E. Known to occur in more than four of the habitats given at A2.3, with at least four a natural habitat. 6
- U. Unknown

Score 6

Documentation:

Identify type of habitats where it occurs and degree/type of impacts:
 See A 2.3.
 Sources of information:
 Tu, 2003; Brooklyn Botanic Garden, 2008.

3.3. Role of disturbance in establishment

- A. Requires anthropogenic disturbances to establish. 0
- B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
- C. Can establish independent of any known natural or anthropogenic disturbances. 4
- U. Unknown

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Score 4

Documentation:

Identify type of disturbance:

Generally found in areas with disturbance in NY metro area, but in the Adirondacks yellow iris rapidly invades undisturbed, pristine areas (e.g. Siamese Wilderness Area; S. Flint 2008 pers. comm.)

Sources of information:

Cody, 1961; Dalton, 1983; Tu, 2003; Gravuer, 2005; Invasive Species Specialist Group 2005; author's (Moore's) personal observations; Steve Glenn's personal observations.

3.4. Climate in native range

- A. Native range does not include climates similar to New York 0
- B. Native range possibly includes climates similar to at least part of New York. 1
- C. Native range includes climates similar to those in New York 3
- U. Unknown

Score 3

Documentation:

Describe what part of the native range is similar in climate to New York:

Range in Europe and Asia with climates similar to NY.

Sources of information:

Gravuer, 2005; Brooklyn Botanic Garden, 2008.

3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope)

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score 4

Documentation:

Identify states and provinces invaded:

CT, DC, DE, IA, IL, IN, MA, MD, ME, MI, MN, NH, NJ, NY, OH, PA, RI, VA, VT, WI, WV; NB, NF, NS, ON, PE, QC.

Sources of information: See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.

U.S.D.A., 2008.

3.6. Current introduced distribution of the species in natural areas in the eight New York State PRISMs (Partnerships for Regional Invasive Species Management)

- A. Present in none of the PRISMs 0
- B. Present in 1 PRISM 1
- C. Present in 2 PRISMs 2
- D. Present in 3 PRISMs 3
- E. Present in more than 3 PRISMs or on the Federal noxious weed lists 4

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

U. Unknown

Score

| |
|---|
| 4 |
|---|

| | |
|--|--|
| Documentation: | |
| Describe distribution: | |
| Present in all PRISMs | |
| Sources of information: | |
| Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008. | |

Total Possible

| |
|----|
| 25 |
|----|

Section Three Total

| |
|----|
| 25 |
|----|

4. DIFFICULTY OF CONTROL

4.1. Seed banks

- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
- B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
- C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
- U. Unknown

Score

| |
|---|
| 2 |
|---|

| | |
|--|--|
| Documentation: | |
| Identify longevity of seed bank: | |
| Evidence that seeds can germinate the season after ripening. | |
| Sources of information: | |
| Dymes, 1920; Sutherland, 1990; Peat & Fitter, 2005. | |

4.2. Vegetative regeneration

- A. No regrowth following removal of aboveground growth 0
- B. Regrowth from ground-level meristems 1
- C. Regrowth from extensive underground system 2
- D. Any plant part is a viable propagule 3
- U. Unknown

Score

| |
|---|
| 2 |
|---|

| | |
|---------------------------------------|--|
| Documentation: | |
| Describe vegetative response: | |
| Extensive underground rhizome system. | |
| Sources of information: | |
| Tu, 2003; Gravuer, 2005. | |

4.3. Level of effort required

- A. Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. 0
- B. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft²). 2
- C. Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). 3
- D. Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. 4

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- U. Unknown
Eradication may be impossible (infestation as above).

Score

| |
|---|
| 4 |
|---|

Documentation:

Identify types of control methods and time-term required:
 Mechanical methods useful for small populations where one can be sure that all of the underground tubers are removed (long-lived seed bank does not exist). However, complete removal by mechanical methods is not possible for larger infestations. Early season use of Rodeo has been shown to be effective (Russ McClain, pers. obs. in Tu, 2003). Stem injection is the most effective method in the Adirondacks, but the seasonal window for treatment is small, thus limiting the number and size of infestations that can be controlled. Note that some people are sensitive to the resinous substances found in the leaves.
 Sources of information:
 Jacono, 2001; Tu, 2003; S. Flint 2008 pers. comm.

Total Possible

| |
|----|
| 10 |
|----|

Section Four Total

| |
|---|
| 8 |
|---|

Total for 4 sections Possible

| |
|-----|
| 100 |
|-----|

Total for 4 sections

| |
|----|
| 76 |
|----|

C. STATUS OF CULTIVARS AND HYBRIDS:

At the present time (May 2008) there is no protocol or criteria for assessing the invasiveness of cultivars independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain, and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Some cultivars of the species known to be available: ‘Alba,’ ‘Gigantea,’ ‘Golden Fleece,’ ‘Holden Clough,’ ‘Mandschurica’

References for species assessment:

Brooklyn Botanic Garden. 2008. AILANTHUS database. [Accessed on 17 March 2008, 1 July 2008.].

Cody, W. J. 1961. *Iris pseudacorus* L. escaped from cultivation in Canada *Canad. Field-Naturalist* 75: 139-142.

Coops, H. and G. Van Der Velde. 1995. Seed dispersal, germination and seedling growth of six helophyte species in relation to water-level zonation. *Freshwater Biology* 34: 13-20.

Cox, G.W. 1999. *Alien species in North America and Hawaii: Impacts on Natural Ecosystems*. Island Press, Washington, D.C.

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

Dalton, P. A. 1983. Aquatic and wetland plants of the Arnold Arboretum. *Arnoldia* (Jamaica Plain) 43: 7-44.

Dymes, T.A. 1920. Notes on the life history of *Iris pseudacorus*, Linn., with special reference to its seeds and seedlings. *Proceedings of the Linnean Society of London* 132: 59-63.

Flint, Steven. 2008. The Nature Conservancy, Terrestrial Project Coordinator, Adirondacks. Rt 73, Keene Valley, NY12943. sflint@tnc.org. Personal communication to Marilyn Jordan.

Gleason, H. A. & A. Cronquist. 1991. *Manual of Vascular Plants of Northeastern United States and Adjacent Canada*. 2nd ed. The New York Botanical Garden, Bronx. 993 pp.

Goldblatt, P. 2002. *Iris*. In: *Flora of North America* Editorial Committee, eds. 1993+. *Flora of North America North of Mexico*. New York and Oxford. Vol. 26, pp. 390.

Gravuer, K. 2005. U.S. Invasive Species Impact Rank (I-Rank) for *Iris pseudacorus*. <natureserve.org> [Accessed on 1 July 2008].

Han, Y. L., S. Qiu, C. Y. Xia, J. Su & S. Z. Huang 2006. Selection of adaptable environment for *Iris pseudacorus* growth *Journal-of-Plant-Resources-and-Environment* 15(2): 38-41.

Heckman, A. 2005. Part IV. Plant Assessment Form, for use with "Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands", Western Washington and Western Oregon. *Iris pseudacorus*. <invasivespeciescoalition.org/GardenPlants/YellowFlagIrisAssessment.pdf> [Accessed 17 March 2008]

Invasive Species Specialist Group 2005. *Global Invasive Species Database* <<http://www.issg.org/database/species/ecology.asp?si=873&fr=1&sts=sss>>. [Accessed 17 March 2008].

Jacono, C.C. 2000, 30 September 2001 last update. *Iris pseudacorus*. United States Department of the Interior, Biological Resources Division, Center for Aquatic Resource Studies. Online. Available: http://nas.er.usgs.gov/taxgroup/plants/docs/ir_pseud.html [Accessed 1 July 2008].

Mehrhoff, L.J., J.A. Silander, Jr., S.A. Leicht and E. Mosher. 2003. IPANE: Invasive Plant Atlas of New England. Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT. Online. Available: <http://invasives.eeb.uconn.edu/ipane/>.

Peat, H., and A. Fitter. 2005. *The Ecological Flora of the British Isles at the University of York*. <www.york.ac.uk/res/ecoflora/cfm/ecofl/index.cfm>. [Accessed 1 July 2008].

Ramey, V. and B. Peichel. 2001. *Non-Native Invasive Aquatic Plants in the United States: Iris pseudacorus*. Center for Aquatic and Invasive Plants, University of Florida and Sea Grant, University of Minnesota. Online. <aquat1.ifas.ufl.edu/seagrant/iripse2.html> [Accessed 1 July 2008].

Randall, R.P. 2002. *A global compendium of weeds*. R.G. and F.J. Richardson, Melbourne. 905 pp.

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

Sutherland, W. J. 1990. *Iris pseudacorus* L. *Journal of Ecology* 78(3): 833-848.

Thomas, L. K. 1980. The impact of three exotic plant species on a Potomac island. National Park Service scientific monograph series; No. 13. U.S. Department of the Interior, Washington, D.C. 179 pp.

Tu, M. 2003. Element stewardship abstract: *Iris pseudacorus*. The Nature Conservancy. Online. Available: <tncweeds.ucdavis.edu/esadocs/documnts/irisps.html>. [Accessed 1 July 2008]

United States Department of Agriculture, National Resources Conservation Service. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana [Accessed on 17 March 2008.].

Washington Invasive Species Coalition (WISC). 2004. Plants invasive to western Washington: Yellow flag iris. <invasivespeciescoalition.org/GardenPlants/InvasivePlants/document_view> [Accessed 1 July 2008.].

Washington State Noxious Weed Control Board. 2003. Written findings of the State Noxious Weed Control Board. <nwcb.wa.gov/weed_info/contents.html>. [Accessed 1 July 2008].

Weber, E. 2003. *Invasive plant species of the world: a reference guide to environmental weeds*. CABI Publishing, Cambridge, Massachusetts. 548 pp.

Whitinger, D. 2005. Dave's Garden: PlantFiles. <davesgarden.com/pdf/>. [Accessed 1 July 2008].

Weldy, T. and D. Werier. 2005. *New York Flora Atlas*. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. [Accessed on 1 July 2008].

Citation: This NY ranking form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

Acknowledgments: The NY form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form. Original members of the LIISMA SRC included representatives of the Brooklyn Botanic Garden; The Nature Conservancy; New York Natural Heritage Program, New York Sea Grant; New York State Office of Parks, Recreation and Historic Preservation; National Park Service; Brookhaven National Laboratory; New York State Department of Environmental Conservation Region 1; Cornell Cooperative Extension of Suffolk/Nassau Counties; Long Island Nursery and Landscape Association; Long Island Farm Bureau; SUNY Farmingdale Ornamental Horticulture Department; Queens College Biology Department; Long Island Botanical Society; Long Island Weed Information Management System database manager; Suffolk County Department of Parks, Recreation and Conservation; Nassau County Department of Parks, Recreation and Museums; Suffolk County Soil & Water Conservation District.

References for ranking form:

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

- Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.
- Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).
- Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>
- Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49
- Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.
- Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.