

# NEW YORK

## TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM

Scientific name: Alopochen aegyptiacus  
 Common names: Egyptian Goose  
 Native distribution: Africa  
 Date assessed: 7/1/2013  
 Assessors: D. Adams  
 Reviewers: \_\_\_\_\_  
 Date Approved: \_\_\_\_\_ Form version date: 3 January 2013

**New York Invasiveness Rank:** Moderate (Relative Maximum Score 50.00-69.99)

<b>Distribution and Invasiveness Rank</b> ( <i>Obtain from PRISM invasiveness ranking form</i> )		
Status of this species in each PRISM:	Current Distribution	PRISM Invasiveness Rank
1 Adirondack Park Invasive Program	Not Present	Not Assessed
2 Capital/Mohawk	Not Present	Not Assessed
3 Catskill Regional Invasive Species Partnership	Not Present	Not Assessed
4 Finger Lakes	Not Present	Not Assessed
5 Long Island Invasive Species Management Area	Not Present	Not Assessed
6 Lower Hudson	Not Present	Not Assessed
7 Saint Lawrence/Eastern Lake Ontario	Not Present	Not Assessed
8 Western New York	Not Present	Not Assessed

<b>Invasiveness Ranking Summary</b> (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	30 ( <u>30</u> )	13
2	Biological characteristic and dispersal ability	30 ( <u>30</u> )	23
3	Ecological amplitude and distribution	30 ( <u>30</u> )	11
4	Difficulty of control	10 ( <u>10</u> )	5
	Outcome score	100 ( <u>100</u> ) <sup>b</sup>	52 <sup>a</sup>
	Relative maximum score †		52
	New York Invasiveness Rank §	Moderate (Relative Maximum Score 50.00-69.99)	

\* For questions answered "unknown" do not include point value in "Total Answered Points Possible." If "Total Answered Points Possible" is less than 70.00 points, then the overall invasive rank should be listed as "Unknown."

† Calculated as 100(a/b) to two decimal places.

§ Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00

### A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

A1.1. Has this species been documented in NY? (reliable source; voucher not required)		
<input type="checkbox"/>	Yes – continue to A1.2	
<input checked="" type="checkbox"/>	No – continue to A2.1; Yes <input checked="" type="checkbox"/> NA; Yes <input checked="" type="checkbox"/> USA	
A1.2. In which PRISMs is it known (see inset map)?		
<input type="checkbox"/>	Adirondack Park Invasive Program	
<input type="checkbox"/>	Capital/Mohawk	
<input type="checkbox"/>	Catskill Regional Invasive Species Partnership	
<input type="checkbox"/>	Finger Lakes	
<input type="checkbox"/>	Long Island Invasive Species Management Area	
<input type="checkbox"/>	Lower Hudson	
<input type="checkbox"/>	Saint Lawrence/Eastern Lake Ontario	

# NEW YORK

## TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM

<input type="checkbox"/>	Western New York
--------------------------	------------------

**Documentation:**

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

A2.0. Is this species listed on the Federal Injurious Fish and Wildlife list?

- Yes – the species will automatically be listed as Prohibited, no further assessment required  
 No – continue to A2.1

A2.1. What is the likelihood that this species will occur and persist given the climate in the following PRISMs?  
 (obtain from PRISM invasiveness ranking form and/ or Climatch score)

- |                   |  |
|-------------------|--|
| Unlikely          | Adirondack Park Invasive Program               |
| Moderately Likely | Capital/Mohawk                                 |
| Unlikely          | Catskill Regional Invasive Species Partnership |
| Moderately Likely | Finger Lakes                                   |
| Very Likely       | Long Island Invasive Species Management Area   |
| Moderately Likely | Lower Hudson                                   |
| Moderately Likely | Saint Lawrence/Eastern Lake Ontario            |
| Moderately Likely | Western New York                               |

**Documentation:**

Sources of information (e.g.: distribution models, literature, expert opinions):  
 Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

***If the species does not occur and is not likely to survive and reproduce within any of the PRISMs, then stop here as there is no need to assess the species.***

A2.2. What is the current distribution of the species in each PRISM? (obtain rank from PRISM invasiveness ranking forms)

Adirondack Park Invasive Program	Distribution
Capital/Mohawk	Not Present
Catskill Regional Invasive Species Partnership	Not Present
Finger Lakes	Not Present
Long Island Invasive Species Management Area	Not Present
Lower Hudson	Not Present
Saint Lawrence/Eastern Lake Ontario	Not Present
Western New York	Not Present

**Documentation:**

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

A2.3. Describe the potential or known suitable habitats within New York. Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

Aquatic Habitats	Wetland Habitats	Upland Habitats
<input type="checkbox"/> Marine	<input type="checkbox"/> Salt/brackish marshes	<input type="checkbox"/> Cultivated*
<input type="checkbox"/> Salt/ brackish waters	<input checked="" type="checkbox"/> Freshwater marshes	<input checked="" type="checkbox"/> Grasslands/old fields
<input type="checkbox"/> Freshwater tidal	<input type="checkbox"/> Peatlands	<input type="checkbox"/> Shrublands
<input type="checkbox"/> Rivers/streams	<input type="checkbox"/> Shrub swamps	<input type="checkbox"/> Forests/woodlands
<input checked="" type="checkbox"/> Natural lakes and ponds	<input type="checkbox"/> Forested wetlands/riparian	<input type="checkbox"/> Alpine
<input type="checkbox"/> Vernal pools	<input type="checkbox"/> Ditches*	<input type="checkbox"/> Roadsides*
<input checked="" type="checkbox"/> Reservoirs/impoundments*	<input type="checkbox"/> Beaches/or coastal dunes	<input checked="" type="checkbox"/> Cultural*

Other potential or known suitable habitats within New York:

Favours a diversity of wetlands in open country, only shunning densely-wooded areas, also occurs on

**NEW YORK**  
**TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

---

meadows and grasslands.

**Documentation:**

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

**B. INVASIVENESS RANKING**

*1. ECOLOGICAL IMPACT*

1.1. Impact on Ecosystem Processes and System-wide Parameters (e.g., energy cycle, nutrient and mineral dynamics, light availability, or geomorphological changes (erosion and sedimentation rates).

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years 0
- B. Influences ecosystem processes to a minor degree, has a perceivable but mild influence 3
- C. Significant alteration of ecosystem processes 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes 10
- U. Unknown

Score

**Documentation:**

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)

Forages mainly by grazing on pasture and arable land, also by dabbling on surface and with head submerged. Droppings may cause eutrophication of still waters.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

1.2. Impact on Natural Habitat

- A. No perceived impact; causes no apparent change in native habitat 0
- B. Influences natural habitat (e.g., reduces the stem density and height of one or more native species in core habitat) 3
- C. Significantly alters natural habitat (e.g., produces a notable reduction in the population size of one or more native species in core habitat) 7
- D. Causes major alteration in natural habitat (e.g., results in the extirpation of one or more native species, or changes the community composition in core habitat towards species exotic to the natural community) 10
- U. Unknown

Score

**Documentation:**

Identify type of impact or alteration:

Feeds on grass and seeds, leaves and stems of plants, vegetables, grains, shoots and potatoes, also some animal items such as worms and locusts. Large groups may cause physical damage to habitats through grazing or trampling.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

1.3. Impact on other species or species groups, including cumulative impact of this species on other organisms in the community it invades (e.g., interferes with native predator/ prey dynamics; hybridizes with a native species; hosts a non-native disease which impacts a native species)

**NEW YORK  
TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

- A. Negligible perceived impact 0
- B. Minor impact (e.g. 1 species, <20% population decline) 3
- C. Moderate impact (e.g. 2-3 species and/ or 20-29% population decline of any 1 species) 7
- D. Severe impact on other species or species groups (e.g. >3 species and/ or ≥30% population decline of any 1 species) 10
- U. Unknown

Score 3

**Documentation:**

Identify type of impact or alteration:

Share feeding habitat with mallards, other ducks and coots and may compete with them for food or territories. Threat to native endemic flora and fauna on islands. Common to locally abundant with greatest numbers in S and E Africa. In parts of range regarded as agricultural pest. Populations have increased in S Africa during the present century due to building of dams and irrigation systems.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

Total Possible 30  
Section One Total 13

**2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY**

**2.1. Mode and rate of reproduction**

- A. No reproduction (e.g. sterile with no sexual or asexual reproduction) 0
- B. Limited reproduction (e.g. intrinsic rate of increase <10%/ year) 1
- C. Moderate reproduction (e.g. intrinsic rate of increase between 10-30%/ year) 2
- D. Abundant reproduction (e.g. intrinsic rate of increase >30%/ year) 4
- U. Unknown

Score 2

**Documentation:**

Describe key reproductive characteristics:

Has high reproductive potential. Gregarious. Long lived. Shared incubation, parental care of young.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

**2.2. Migratory behavior**

- A. Always migratory in its native range 0
- B. Non-migratory or facultative migrant in its native range 2
- U. Unknown

Score 2

**Documentation:**

Describe migratory behavior:

Highly mobile locally. May migrate northward from the Nile Valley in its native range.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

**2.3. Biological potential for colonization by long-distance dispersal/ movement.**

- A. No long-distance dispersal/ movement mechanisms 0
- A. Adaptations exist for long-distance dispersal, but studies report that most individuals (90%) establish territories within 10 miles of parent or within a distance twice the home range of 1

**NEW YORK  
TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

---

- the parent, and tend not to cross major barriers such as rivers and major roads
- B. Adaptations exist for long-distance dispersal, movement and evidence that offspring often disperse greater than 10 miles of parent or greater than twice the home range of parent and will cross major barriers such as river and major roads 2
- U. Unknown

Score

**Documentation:**

Identify dispersal mechanisms:

While not breeding, it disperses somewhat, sometimes making longer migrations northward from the Nile Valley.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

2.4. Practical potential to be spread by human activities, both directly and indirectly (possible vectors include: commercial sales, deliberate stocking, translocation, rehabilitation, pest control industry, agricultural escapes, pet abandonment and release, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 4
- U. Unknown

Score

**Documentation:**

Identify dispersal mechanisms:

Benefits from human association. Introduced to Britain in the 17th century, feral populations exist including Florida and New Zealand.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

2.5. Non-living chemical and physical characteristics that increase competitive advantage (e.g., tolerance to various extremes, pH, temperature, fill vacant niche, charismatic species)

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 4
- C. Possesses two or more characteristics that increase competitive advantage 8
- U. Unknown

Score

**Documentation:**

Evidence of competitive ability:

Raised for food and extensively bred in parts of Africa since domesticated by the Egyptians. Small feral populations have become established in western Europe from chiefly ornamental birds. Will nest in a wide variety of situations.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

2.6. Biological characteristics that increase competitive advantage (e.g., high fecundity, generalist, highly evolved defense mechanisms, behavioral adaptations)

**NEW YORK  
TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

- |    |   |   |
|----|---|---|
| A. | Possesses no characteristics that increase competitive advantage          | 0 |
| B. | Possesses one characteristic that increases competitive advantage         | 4 |
| C. | Possesses two or more characteristics that increase competitive advantage | 8 |
| U. | Unknown   |   |

Score 8

**Documentation:**  
 Evidence of competitive ability:  
 Aggressively territorial towards their own species when breeding. Both parents take turns incubating eggs. Pair for life. Care for young until they are old enough to care for themselves.  
 Sources of information:  
 Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
 & Invasive Species Compendium

- 2.7. Other species in the family and/ or genus invasive in New York or elsewhere?
- |    |         |   |
|----|---------|---|
| A. | No      | 0 |
| B. | Yes     | 2 |
| U. | Unknown |   |

Score 0

**Documentation:**  
 Identify species:

	Total Possible	<span style="border: 1px solid black; padding: 2px 10px;">30</span>
	Section Two Total	<span style="border: 1px solid black; padding: 2px 10px;">23</span>

**3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION**

- 3.1. Current introduced distribution of established populations in the northern latitudes of USA and southern latitude of Canada (e.g., between 35 and 55 degrees)
- |    |   |   |
|----|---|---|
| A. | Not known from the northern US or southern Canada   | 0 |
| B. | Established as a non-native in 1 northern USA state and/or southern Canadian province   | 1 |
| C. | Established as a non-native in 2 or 3 northern USA states and/or southern Canadian provinces  | 2 |
| D. | Established as a non-native in 4 or more northern USA states and/or southern Canadian provinces, and/or categorized as a problem species (e.g., “Invasive”) in 1 northern state or southern Canadian province | 3 |
| U. | Unknown   |   |

Score 0

**Documentation:**  
 Identify states and provinces:  
 Current populations in US limited to FL.  
 Sources of information:  

- See known introduced range at [www.usda.gov](http://www.usda.gov), and update with information from states and Canadian provinces.

 Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
 & Invasive Species Compendium

- 3.2. Current introduced distribution of established populations of the species in natural areas in the eight New York State PRISMs (Partnerships for Regional Invasive Species Management)
- |    |                                   |   |
|----|-----------------------------------|---|
| A. | Established in none of the PRISMs | 0 |
|----|-----------------------------------|---|

**NEW YORK**  
**TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

---

- B. Established in 1 PRISM 1
- C. Established in 2 or 3 PRISMs 3
- D. Established in 4 or more PRISMs 5
- U. Unknown

Score

**Documentation:**

Describe distribution:

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

3.3. Number of known, or potential (each individual possessed by a vendor or consumer is a potential release), individual releases and/ or release events (propagule pressure)

- A. None 0
- B. Few releases (e.g., <10 annually) 2
- C. Regular, small scale releases (e.g., 10-99 annually) 4
- D. Multiple, large scale (e.g.,  $\geq 100$  annually) 6
- U. Unknown

Score

**Documentation:**

Describe known or potential releases:

Widely domesticated, ornamental species.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

3.4. Current introduced population density in northern USA and/ or southern Canada.

- A. No known populations established 0
- B. Low to moderate population density (e.g.,  $\leq 1/4$  or  $<$  to  $1/2$  native population density) 1
- C. High or irruptive population density (e.g.,  $\geq 1/2$  native population density) 2
- U. Unknown

Score

**Documentation:**

Describe population density:

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

3.5. Number of habitats the species may invade

- A. Not known to invade any natural habitats 0
- B. Known to occur in 2/ 3 habitats, with at least 1/ 2 natural habitat(s) 2
- C. Known to occur in 4 or more habitats, with at least 3 natural habitats 3
- U. Unknown

Score

**Documentation:**

**NEW YORK  
TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

Identify type of habitats where it occurs and degree/type of impacts:

Favors a diversity of wetlands in open country.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

3.6. Role of anthropogenic (human related) features in establishment (e.g. buildings, roads, agricultural fields, etc)

- |    |   |   |
|----|---|---|
| A. | Requires anthropogenic disturbances to establish  | 0 |
| B. | May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances | 2 |
| C. | Can establish independent of any known natural or anthropogenic disturbances  | 3 |
| U. | Unknown   |   |

Score 3

**Documentation:**

Identify anthropogenic features:

Benefits from cultivation. In parts of its range considered an agricultural pest.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

3.7. Climate in native range (e.g., med. to high,  $\geq 5$ , Climatch score; within 35 to 55 degree latitude; etc.)

- |    |   |   |
|----|---|---|
| A. | Native range does not include climates similar to New York              | 0 |
| B. | Native range possibly includes climates similar to portions of New York | 4 |
| C. | Native range includes climates similar to those in New York             | 8 |
| U. | Unknown   |   |

Score 4

**Documentation:**

Describe known climate similarities: 54 degrees N to 34 degrees S. Established in Britain.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

Total Possible	30
Section Three Total	11

**4. DIFFICULTY OF CONTROL**

4.1. Re-establishment potential, nearby propagule source, known vectors of re-introduction in vicinity (e.g. biological supplies, pets, game farms, zoos, shooting preserves, connecting corridors, mechanized transportation)

- |    |   |   |
|----|---|---|
| A. | No known vectors/ propagule source for re-establishment following removal                         | 0 |
| B. | Possible re-establishment from 1 vector/ propagule source following removal                       | 1 |
| C. | Likely to re-establish from 2-3 vectors/ propagule sources following removal                      | 2 |
| D. | Strong potential for re-establishment from 4 or more vectors/ propagule sources following removal | 3 |
| U. | Unknown   |   |

Score 1

**Documentation:**

Identify source/ vectors:

Widely domesticated, ornamental species.

Sources of information:



**NEW YORK  
TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

4.2. Status of monitoring and/ or management protocols for species

- A. Standardized protocols appropriate to New York State are available 0
- B. Scientific protocols are available from other countries, regions or states 1
- C. No known protocols exist 2
- U. Unknown

Score

**Documentation:**

Describe protocols:

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

4.3. Status of monitoring and/ or management resources (e.g. tools, manpower, travel, traps, lures, ID keys, taxonomic specialists, etc.)

- A. Established resources are available including commercial and/ or research tools 0
- B. Monitoring resources may be available (e.g. partnerships, NGOs, etc) 1
- C. No known monitoring resources are available 2
- U. Unknown

Score

**Documentation:**

Describe resources:

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

4.4. Level of effort required.

- A. Management is not required (e.g., species does not persist without repeated human mediated action) 0
- B. Management is relatively easy and inexpensive; invasive species can be maintained at low abundance causing little or no ecological harm (e.g., 10 or fewer person-hours of manual effort can eradicate a local population in 1 year) 1
- C. Management requires a major short-term investment, and is logistically and politically challenging; eradication is difficult, but possible (e.g., 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/ year for 2-5 years to suppress a local population) 2
- D. Management requires a major investment and is logistically and politically difficult; eradication may be impossible (e.g., more than 100 person-hours/ year of manual effort, or more than 10 person hours/year for more than 5 years to suppress a local population) 3
- U. Unknown

Score

**Documentation:**

Identify types of control methods and time required:

Populations in the US currently limited to Florida.

Sources of information:

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.  
& Invasive Species Compendium

Total Possible

Section Four Total

**Total for 4 sections Possible**

**NEW YORK  
TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

---

**Total for 4 sections**

52
----

**C. STATUS OF GENETIC VARIANTS AND HYBRIDS:**

At the present time there is no protocol or criteria for assessing the invasiveness of genetic variants independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

Genetic variants of the species known to exist: *A. aegyptiacus* is the only extant member of the genus *Alopochen*.

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain, and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Hybrids of uncertain origin known to exist: Hybridization has occurred between *A. aegyptiaca* and the Orinoco goose *Neochen jubata* in captivity, and apparent hybrids with mallard *Anas platyrhynchos*, ruddy shelduck *Tadorna ferruginea*, Canada goose *Branta canadensis* and domestic geese.

**References for species assessment:**

Handbook of the Birds of the World. 1992. Editors: Hyo, Elliot and Sargatal.

Invasive Species Compendium. [www.cabi.org/isc/](http://www.cabi.org/isc/)

**Citation:** The New York Terrestrial Vertebrate Invasiveness Ranking Form is an adaptation of the New York Plant Invasiveness Ranking Form. The original plant form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY.

**Acknowledgments:** The New York Terrestrial Vertebrate Invasiveness Ranking Form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Invasive Species Council and Invasive Species Advisory Committee were incorporated in revisions of this form. Members of the Office of Invasive Species Coordination's Four-tier Team, who coordinated the effort, included representatives of the New York State Department of Environmental Conservation\* (Division of Fish, Wildlife and Marine Resources, Division of Lands and Forests, Division of Water); The Nature Conservancy; New York Natural Heritage Program; New York Sea Grant; Lake Champlain Sea Grant\*; New York State Department of Agriculture and Markets (Division of

**NEW YORK**  
**TERRESTRIAL VERTEBRATE INVASIVENESS RANKING FORM**

---

Plant Industry and Division of Animal Industry); Cornell University (Department of Natural Resources and Department of Entomology); New York State Nursery and Landscape Association; New York Farm Bureau; Brooklyn Botanic Garden; Pet Industry Joint Advisory Council\*; Trout Unlimited; United States Department of Agriculture Animal and Plant Health Inspection Service\* (Plant Protection and Quarantine and Wildlife Services); New York State Department of Transportation; State University of New York at Albany and Plattsburgh; and Cary Institute of Ecosystem Studies. Those organizations listed with an asterisk comprised the Terrestrial Vertebrate Working Group.

**References for ranking form:**

Bomford, M. 2008. Risk Assessment Models for Establishment of Exotic Vertebrates in Australia and New Zealand. Invasive Animals Cooperative Research Centre, Canberra.

Broken Screens: The Regulation of Live Animal Imports in the United States. 2007. Defenders of Wildlife, Washington, DC.

Cooperative Prevention of Invasive Wildlife Introduction in Florida. 2008. The Environmental Law Institute, Washington, DC.

Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, New York.

Natural Resources Board Order No. IS-34-06, Invasive Species Identification, Classification and Control. 2008. Wisconsin Department of Natural Resources, Madison Wisconsin.

Preventing Biological Invasions: Best Practices in Pre-Import Risk Screening for Species of Live Animals in International Trade. 2008. Convention of Biological Diversity, Global Invasive Species Programme and Invasive Species Specialist Group of IUCN's Species Survival Commission. University of Notre Dame, Indiana.

Standard Methodology to Assess the Risks From Non-native Species Considered Possible Problems to the Environment. 2005. DEFRA.

Witmer, G., W. Pitt and K. Fagerstone. 2007. Managing Vertebrate Invasive Species. USDA National Wildlife Research Center Symposia, Fort Collins, Colorado.