

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Scientific name: Pseudosasa japonica (Siebold & Zucc. ex Steud.) Makino ex Nakai
 USDA Plants Code: PSJA

Common names: Arrow bamboo

Native distribution: Eastern Asia

Date assessed: February 21, 2010; revised January 17, 2013

Assessors: Gerry Moore; revised by Marilyn Jordan and SRC

Reviewers: LIISMA SRC

Date Approved: March 10, 2010; ___2013 Form version date: 28 November 2012

New York Invasiveness Rank: Not Assessable

Distribution and Invasiveness Rank (Obtain from PRISM invasiveness ranking form)			
	Status of this species in each PRISM:	Current Distribution	PRISM Invasiveness Rank
1	Adirondack Park Invasive Program	Not Assessed	Not Assessed
2	Capital/Mohawk	Not Assessed	Not Assessed
3	Catskill Regional Invasive Species Partnership	Not Assessed	Not Assessed
4	Finger Lakes	Not Assessed	Not Assessed
5	Long Island Invasive Species Management Area	Not Present	Moderate
6	Lower Hudson	Not Assessed	Not Assessed
7	Saint Lawrence/Eastern Lake Ontario	Not Assessed	Not Assessed
8	Western New York	Not Assessed	Not Assessed

Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 (30)	17
2	Biological characteristic and dispersal ability	25 (22)	11
3	Ecological amplitude and distribution	25 (25)	14
4	Difficulty of control	10 (10)	6
	Outcome score	100 (87) ^b	48 ^a
	Relative maximum score † If assessable (escapes cultivation) would rank M		(55.17)
	New York Invasiveness Rank (for natural areas) [§]	Not Assessable	

* For questions answered “unknown” do not include point value in “Total Answered Points Possible.” If “Total Answered Points Possible” is less than 70.00 points, then the overall invasive rank should be listed as “Unknown.”

† Calculated as 100(a/b) to two decimal places.

§ Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00

Not Assessable: not persistent in NY, or not found outside of cultivation.

A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

A1.1. Has this species been documented to persist without cultivation in NY? (reliable source; voucher not required)		
<input type="checkbox"/>	Yes – continue to A1.2	
<input checked="" type="checkbox"/>	No – continue to A2.1	
A1.2. In which PRISMs is it known (see inset map)?		
<input type="checkbox"/>	Adirondack Park Invasive Program	
<input type="checkbox"/>	Capital/Mohawk	
<input type="checkbox"/>	Catskill Regional Invasive Species Partnership	
<input type="checkbox"/>	Finger Lakes	
<input type="checkbox"/>	Long Island Invasive Species Management Area	
<input type="checkbox"/>	Lower Hudson	
<input type="checkbox"/>	Saint Lawrence/Eastern Lake Ontario	
<input type="checkbox"/>	Western New York	

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Documentation: Weldy & Werier (2010) do not report *Pseudosasa japonica* as an escape in New York. Population in Alley Pond Park (Queens Co.) is probably spreading vegetatively from plantings. The species is also planted at the Brooklyn Botanic Garden but no natural escapes from cultivation are known in NYS. Moore(2013) has noted this species in New Jersey at one site in Burlington Co. One occurrence in Westport CT occurs in "shrubby waste" and probably resulted from human actions.

Sources of information:

Stapleton, 2007a; Brooklyn Botanic Garden, 2023; Weldy & Werier, 2010; U.S.D.A., 2010.

A2.1. What is the likelihood that this species will occur and persist outside of cultivation, given the climate in the following PRISMs? (obtain from PRISM invasiveness ranking form)

Not Assessed	Adirondack Park Invasive Program
Not Assessed	Capital/Mohawk
Not Assessed	Catskill Regional Invasive Species Partnership
Not Assessed	Finger Lakes
Unlikely	Long Island Invasive Species Management Area
Not Assessed	Lower Hudson
Not Assessed	Saint Lawrence/Eastern Lake Ontario
Not Assessed	Western New York

Documentation: Since the species is unlikely to produce viable seeds and escape cultivation by sexual or vegetative propagules its rank was changed from Moderate to Not Assessable when this assessment was revised in 2013. See Question 2.1 for more information about flowering and vegetative growth. NYS point score is retained to indicate invasive potential in natural or cultivated areas should the species escape cultivation in the future.

Sources of information (e.g.: distribution models, literature, expert opinions):

Brooklyn Botanic Garden, 2013. Stapleton, 2007b; Moore 2010 pers. obs.; SRC obs.

If the species does not occur and is not likely to occur in any of the PRISMs, then stop here as there is no need to assess the species. Rank is "Not Assessable."

A2.2. What is the current distribution of the species in each PRISM? (obtain rank from PRISM invasiveness ranking forms)

	Distribution
Adirondack Park Invasive Program	Not Assessed
Capital/Mohawk	Not Assessed
Catskill Regional Invasive Species Partnership	Not Assessed
Finger Lakes	Not Assessed
Long Island Invasive Species Management Area	Not Present
Lower Hudson	Not Assessed
Saint Lawrence/Eastern Lake Ontario	Not Assessed
Western New York	Not Assessed

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2012; Weldy & Werier NYFA; USDA NRCS.

A2.3. Describe the potential or known suitable habitats within New York. Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

<p>Aquatic Habitats</p> <input type="checkbox"/> Salt/brackish waters <input type="checkbox"/> Freshwater tidal <input type="checkbox"/> Rivers/streams <input type="checkbox"/> Natural lakes and ponds <input type="checkbox"/> Vernal pools <input type="checkbox"/> Reservoirs/impoundments*	<p>Wetland Habitats</p> <input type="checkbox"/> Salt/brackish marshes <input type="checkbox"/> Freshwater marshes <input type="checkbox"/> Peatlands <input type="checkbox"/> Shrub swamps <input checked="" type="checkbox"/> Forested wetlands/riparian <input checked="" type="checkbox"/> Ditches* <input type="checkbox"/> Beaches and/or coastal dunes	<p>Upland Habitats</p> <input checked="" type="checkbox"/> Cultivated* <input checked="" type="checkbox"/> Grasslands/old fields <input type="checkbox"/> Shrublands <input checked="" type="checkbox"/> Forests/woodlands <input type="checkbox"/> Alpine <input checked="" type="checkbox"/> Roadsides*
--	--	---

Other potential or known suitable habitats within New York:

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2010; author's (Moore) pers. obs.

B. INVASIVENESS RANKING

Questions apply to areas similar in climate and habitats to New York unless specified otherwise.

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies OR there are no reports of impacts and the species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score 3

Documentation:

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)

Dense stands clearly reduce light availability, but further studies on impacts to ecosystem processes not known.

Sources of information:

Author's (Moore) pers. comm. and obs.

1.2. Impact on Natural Community Structure

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score 7

Documentation:

Identify type of impact or alteration:

Can form solid dense patches with little to nothing growing below it.

Sources of information:

Author's (Moore) pers. obs.

1.3. Impact on Natural Community Composition

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several) 10

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

native species, reducing biodiversity or change the community composition towards species exotic to the natural community)

U. Unknown

Score

7

Documentation:

Identify type of impact or alteration:

Observed to form a monoculture, significantly reducing the number of native species growing in the community.

Sources of information:

Author's pers. obs.

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades. Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- | | | |
|----|--|----|
| A. | Negligible perceived impact | 0 |
| B. | Minor impact | 3 |
| C. | Moderate impact | 7 |
| D. | Severe impact on other species or species groups | 10 |
| U. | Unknown | |

Score

U

Documentation:

Identify type of impact or alteration:

Impacts on other species or species groups not known.

Sources of information:

Author's (Moore) pers. comm.

	Total Possible	<table border="1" style="display: inline-table;"><tr><td style="width: 20px; height: 20px; text-align: center;">30</td></tr></table>	30
30			
	Section One Total	<table border="1" style="display: inline-table;"><tr><td style="width: 20px; height: 20px; text-align: center;">17</td></tr></table>	17
17			

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction

- | | | |
|----|---|---|
| A. | No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). Such a species should be ranked "Not Assessable" as it will occur only in cultivated settings and cannot escape into natural/minimally managed areas. End the assessment here. | 0 |
| B. | Limited reproduction (fewer than 10 viable seeds per plant; if seed viability is not known, then maximum seed production is less than 100 seeds per plant) AND no reproduction by vegetative propagules (e.g. bulbils, turions, pieces of rhizomes, etc.) is documented as a natural (not spread by people) mode of dispersal across gaps by the species. | 1 |
| C. | Moderate reproduction (fewer than 100 viable seeds per plant; if viability is not known, then maximum seed production is less than 1000 seeds per plant) – OR limited reproduction by vegetative propagules (e.g. bulbils, turions, pieces of rhizomes, etc.) is documented as a natural (not spread by people) mode of dispersal across gaps by the species. For aquatic species viable plant fragments may be treated as vegetative propagules. | 2 |
| D. | Significant reproduction by seeds (more than 100 viable seeds per plant; if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant) –OR abundant reproduction by vegetative propagules (e.g. bulbils, turions, pieces of rhizomes, etc.) is documented as a natural (not spread by people) mode of dispersal across gaps by the species. For aquatic species viable plant fragments may be treated as vegetative propagules. | 4 |
| U. | Unknown | |

Score

0

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Documentation:

Describe key reproductive characteristics (including seeds per plant):

Production of flowers and seeds in native locations is rare, and production of seeds in non-native stands is not known. Plants can spread vegetatively well beyond original planting locations but no conclusive evidence of natural escapes from cultivation could be found (establishment of new populations remote from a planted location without the aid of people). Fragments of rhizomes could be transported by floodwaters but there is no conclusive proof that this has occurred. All known occurrences appear to be either remnants of past cultivation or vegetative spread from a planted specimen and it appears unlikely that this would change in the future. Therefore *P. japonica* scores 0 points for this question and must be ranked "Not Assessable".

Sources of information:

Stapleton, 2007b; author's (Moore) pers. obs.; BBG 2012; SRC observations.

- 2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)
- | | | |
|----|--|---|
| A. | Does not occur (no long-distance dispersal mechanisms) | 0 |
| B. | Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) | 1 |
| C. | Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) | 2 |
| D. | Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) | 4 |
| U. | Unknown | |

Score 1

Documentation:

Identify dispersal mechanisms:

Seed production rare and without obvious adaptations for long distance dispersal. Long distance dispersal can rarely occur when pieces of the rhizomes or stems are moved.

Sources of information:

Stapleton, 2007.

- 2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)
- | | | |
|----|---|---|
| A. | Does not occur | 0 |
| B. | Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) | 1 |
| C. | Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) | 2 |
| D. | High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) | 3 |
| U. | Unknown | |

Score 2

Documentation:

Identify dispersal mechanisms:

Pseudosasa occasionally grown and available commercially. Could be spread by direct means when it is grown or when soils with viable rhizomes or stems present are moved.

Sources of information:

Stapleton, 2007; author's (Moore) pers. comm.

- 2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.
- | | | |
|----|---|---|
| A. | Possesses no characteristics that increase competitive advantage | 0 |
| B. | Possesses one characteristic that increases competitive advantage | 3 |
| C. | Possesses two or more characteristics that increase competitive advantage | 6 |

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS**

U. Unknown

Score 6

Documentation:

Evidence of competitive ability:

Perennial, some shade tolerance, ability to grow on nutrient poor soils; fast growth.

Sources of information:

Clark & Triplett, 2007; author's (Moore) pers. obs.

2.5. Growth vigor

A. Does not form thickets or have a climbing or smothering growth habit 0

B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2

U. Unknown

Score 2

Documentation:

Describe growth form:

Forms a dense layer above shorter vegetation.

Sources of information:

Author's (Moore) pers. obs.

2.6. Germination/Regeneration

A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0

B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2

C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3

U. Unknown (No studies have been completed)

Score U

Documentation:

Describe germination requirements:

Regeneration requirements of rhizomes or stems not known for wild material.

Sources of information:

Author's (Moore) pers. comm.

2.7. Other species in the genus invasive in New York or elsewhere

A. No 0

B. Yes 3

U. Unknown

Score 0

Documentation:

Species:

No other species treated and it is not tracked as invasive. Brooklyn Botanic Garden, 2010; Weldy & Werier, 2010; U.S.D.A. NRCS, 2010.

Total Possible 22
Section Two Total 11

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist: "The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario south of the 47th parallel of latitude")

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score 4

Documentation:
 Identify reason for selection, or evidence of weedy history:
 Stand known from New Jersey was well over a quarter an acre and growing with few other invasives present.
 Sources of information:
 Brooklyn Botanic Garden, 2010; Author's (Moore) pers. obs.

- 3.2. Number of habitats the species may invade**
- A. Not known to invade any natural habitats given at A2.3 0
 - B. Known to occur in one natural habitat given at A2.3 1
 - C. Known to occur in two natural habitats given at A2.3 2
 - D. Known to occur in three natural habitat given at A2.3 4
 - E. Known to occur in four or more natural habitats given at A2.3 6
 - U. Unknown

Score 4

Documentation:
 Identify type of habitats where it occurs and degree/type of impacts:
 See A2.3.
 Sources of information:
 Brooklyn Botanic Garden, 2010; author's (Moore) pers. obs.

- 3.3. Role of disturbance in establishment**
- A. Requires anthropogenic disturbances to establish. 0
 - B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
 - C. Can establish independent of any known natural or anthropogenic disturbances. 4
 - U. Unknown

Score 0

Documentation:
 Identify type of disturbance:
 Generally becomes established only with anthropogenic disturbance.
 Sources of information:
 Brooklyn Botanic Garden, 2010; author's (Moore) pers. obs.

- 3.4. Climate in native range**
- A. Native range does not include climates similar to New York 0
 - B. Native range possibly includes climates similar to at least part of New York. 1
 - C. Native range includes climates similar to those in New York 3
 - U. Unknown

Score 3

Documentation:
 Describe what part of the native range is similar in climate to New York:
 Temperate Asia; established in North America as far north as British Columbia
 Sources of information:

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Stapleton, 2007a; Brooklyn Botanic Garden, 2012; U.S.D.A., 2010

- 3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope)
- A. Not known from the northeastern US and adjacent Canada 0
 - B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
 - C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
 - D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
 - E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces. and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
 - U. Unknown

Score 3

Documentation:

Identify states and provinces invaded:

CT, MD, NJ, PA, VA.

Sources of information: See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.

U.S.D.A. NRCS, 2010.

3.6. Current introduced distribution of the species in natural areas in the eight New York State PRISMs (Partnerships for Regional Invasive Species Management)

- A. Present in none of the PRISMs 0
- B. Present in 1 PRISM 1
- C. Present in 2 PRISMs 2
- D. Present in 3 PRISMs 3
- E. Present in more than 3 PRISMs or on the Federal noxious weed lists 4
- U. Unknown

Score 0

Documentation:

Describe distribution:

See A1.1.

Sources of information:

Brooklyn Botanic Garden, 2010; Weldy & Werier, 2010; U.S.D.A., 2010.

Total Possible 25
Section Three Total 14

4. DIFFICULTY OF CONTROL

- 4.1. Seed banks
- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
 - B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
 - C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
 - U. Unknown

Score 0

Documentation:

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Identify longevity of seed bank:
 Doubtful that rhizomes or stems could remain viable for more than a year. Seed production not known for non-native populations and infrequent in native populations; thus, length of viability not relevant.
 Sources of information:
 Author's (Moore) pers. comm.

- 4.2. Vegetative regeneration
- A. No regrowth following removal of aboveground growth 0
 - B. Regrowth from ground-level meristems 1
 - C. Regrowth from extensive underground system 2
 - D. Any plant part is a viable propagule 3
 - U. Unknown
- Score

2

Documentation:
 Describe vegetative response:
 Extensive underground root system.
 Sources of information:
 Stapleton, 2007a; author's (Moore) pers. obs.

- 4.3. Level of effort required
- A. Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. 0
 - B. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft²). 2
 - C. Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). 3
 - D. Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). 4
 - U. Unknown
- Score

4

Documentation:
 Identify types of control methods and time-term required:
 Species forms large stands including in wetlands, thus requiring a major investment for eradication.
 Removal in wetlands would require a permit.
 Sources of information:
 Author's (Moore) pers. comm.

Total Possible

10

 Section Four Total

6

Total for 4 sections Possible

87

Total for 4 sections

48

C. STATUS OF CULTIVARS AND HYBRIDS:

At the present time (May 2008) there is no protocol or criteria for assessing the invasiveness of cultivars independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain, and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Some cultivars of the species known to be available: 'Tsutsumiana'

References for species assessment:

Brooklyn Botanic Garden. 2013. AILANTHUS database. [Accessed January 2013].

Stapleton, M.A.. 2007a. Pseudosasa. Pp. 27-29 in (Barkworth et al. eds.) Flora North America Vol 24. Oxford University Press, New York. 908 pp.

Stapleton, M.A. 2007b. Bambuseae. Pp. 15-16 in (Barkworth et al. eds.) Flora North America Vol 24. Oxford University Press, New York. 908 pp.

United States Department of Agriculture, National Resources Conservation Service. 2010. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana. < <http://plants.usda.gov/> > [Accessed].

Weldy, T. and D. Werier. 2010. New York Flora Atlas. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. <www.newyork.plantatlas.usf.edu> [Accessed February 21, 2010.].

Citation: This NY ranking form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

Acknowledgments: The NY form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form. Original members of the LIISMA SRC included representatives of the Brooklyn Botanic Garden; The Nature Conservancy; New York Natural Heritage Program, New York Sea Grant; New York State Office of Parks, Recreation and Historic Preservation; National Park Service; Brookhaven National Laboratory; New York State Department of Environmental Conservation Region 1; Cornell Cooperative Extension of Suffolk/Nassau Counties; Long Island Nursery and Landscape Association; Long Island Farm Bureau; SUNY Farmingdale Ornamental Horticulture Department; Queens College Biology Department; Long Island Botanical Society; Long Island Weed Information Management System database manager; Suffolk County Department of Parks, Recreation and Conservation; Nassau County Department of Parks, Recreation and Museums; Suffolk County Soil & Water Conservation District.

References for ranking form:

Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM
FOR NATURAL / MINIMALLY MANAGED AREAS

Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).

Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>

Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49

Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.

Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.