

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

Scientific name:	Acer platanoides	USDA Plants Code: ACPL
Common names:	Norway maple	
Native distribution:	Europe, West Asia (including West Siberia of Russia), and Northern Iran	
Date assessed:	February 26, 2008; September 2, 2008	
Assessors:	Jinshuang Ma, Gerry Moore	
Reviewers:	LIISMA SRC	
Date Approved:	2008-9-10	Form version date: 25 August 2008

New York Invasiveness Rank: Very High (Relative Maximum Score >80.00)

Distribution and Invasiveness Rank (<i>Obtain from PRISM invasiveness ranking form</i>)		
	Status of this species in each PRISM:	PRISM Invasiveness Rank
1	Adirondack Park Invasive Program	Not Assessed
2	Capital/Mohawk	Not Assessed
3	Catskill Regional Invasive Species Partnership	Not Assessed
4	Finger Lakes	Not Assessed
5	Long Island Invasive Species Management Area	High
6	Lower Hudson	Not Assessed
7	Saint Lawrence/Eastern Lake Ontario	Not Assessed
8	Western New York	Not Assessed


Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 (40)	37
2	Biological characteristic and dispersal ability	25 (25)	21
3	Ecological amplitude and distribution	25 (25)	17
4	Difficulty of control	10 (10)	7
	Outcome score	100 (100) ^b	82 ^a
	Relative maximum score †		82.00
	New York Invasiveness Rank §	Very High (Relative Maximum Score >80.00)	

* For questions answered “unknown” do not include point value in “Total Answered Points Possible.” If “Total Answered Points Possible” is less than 70.00 points, then the overall invasive rank should be listed as “Unknown.”

† Calculated as 100(a/b) to two decimal places.

§ Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00

A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

<p>A1.1. Has this species been documented to persist without cultivation in NY? (reliable source; voucher not required)</p> <p><input checked="" type="checkbox"/> Yes – continue to A1.2</p> <p><input type="checkbox"/> No – continue to A2.1</p> <p>A1.2. In which PRISMs is it known (see inset map)?</p> <p><input checked="" type="checkbox"/> Adirondack Park Invasive Program</p> <p><input checked="" type="checkbox"/> Capital/Mohawk</p> <p><input checked="" type="checkbox"/> Catskill Regional Invasive Species Partnership</p> <p><input checked="" type="checkbox"/> Finger Lakes</p> <p><input checked="" type="checkbox"/> Long Island Invasive Species Management Area</p> <p><input checked="" type="checkbox"/> Lower Hudson</p> <p><input checked="" type="checkbox"/> Saint Lawrence/Eastern Lake Ontario</p>	 <p style="font-size: small;">Partnerships for Regional Invasive Species Management 2008</p>
--	--

NEW YORK

NON-NATIVE PLANT INVASIVENESS RANKING FORM

<input checked="" type="checkbox"/>	Western New York
-------------------------------------	------------------

Documentation:

Sources of information:

Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

A2.1. What is the likelihood that this species will occur and persist given the climate in the following PRISMs? (obtain from PRISM invasiveness ranking form)

Not Assessed	Adirondack Park Invasive Program
Not Assessed	Capital/Mohawk
Not Assessed	Catskill Regional Invasive Species Partnership
Not Assessed	Finger Lakes
Very Likely	Long Island Invasive Species Management Area
Not Assessed	Lower Hudson
Not Assessed	Saint Lawrence/Eastern Lake Ontario
Not Assessed	Western New York

Documentation:

Sources of information (e.g.: distribution models, literature, expert opinions):

Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

If the species does not occur and is not likely to occur with any of the PRISMs, then stop here as there is no need to assess the species.

A2.2. What is the current distribution of the species in each PRISM? (obtain rank from PRISM invasiveness ranking forms)

	Distribution
Adirondack Park Invasive Program	Not Assessed
Capital/Mohawk	Not Assessed
Catskill Regional Invasive Species Partnership	Not Assessed
Finger Lakes	Not Assessed
Long Island Invasive Species Management Area	Widespread
Lower Hudson	Not Assessed
Saint Lawrence/Eastern Lake Ontario	Not Assessed
Western New York	Not Assessed

Documentation:

Sources of information:

Fang, 2005; Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

A2.3. Describe the potential or known suitable habitats within New York. Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

Aquatic Habitats	Wetland Habitats	Upland Habitats
<input type="checkbox"/> Salt/brackish waters	<input type="checkbox"/> Salt/brackish marshes	<input checked="" type="checkbox"/> Cultivated*
<input type="checkbox"/> Freshwater tidal	<input type="checkbox"/> Freshwater marshes	<input type="checkbox"/> Grasslands/old fields
<input type="checkbox"/> Rivers/streams	<input type="checkbox"/> Peatlands	<input type="checkbox"/> Shrublands
<input type="checkbox"/> Natural lakes and ponds	<input type="checkbox"/> Shrub swamps	<input checked="" type="checkbox"/> Forests/woodlands
<input type="checkbox"/> Vernal pools	<input checked="" type="checkbox"/> Forested wetlands/riparian	<input type="checkbox"/> Alpine
<input type="checkbox"/> Reservoirs/impoundments*	<input type="checkbox"/> Ditches*	<input checked="" type="checkbox"/> Roadsides*
	<input type="checkbox"/> Beaches and/or coastal dunes	

Other potential or known suitable habitats within New York:

Vacant lots

Documentation:

Sources of information:

Webb & Kaunzinger, 1993; Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

B. INVASIVENESS RANKING

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score 7

Documentation:

Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)
 Good evidence trees cast heavier shade than native tree species (e.g., *Acer rubrum*, oaks) thus significantly decreasing light availability. It has been reported that it has been planted to reduce erosion in areas that are prone to erosion (Love, 2003), but not clear how that should be scored. In contrast, Donnelly suggests that erosion would be worse under Norway Maple because of bare soil beneath the stand, thus leading to water quality problems. Fang (unpub.) measured higher rates of nitrification in soil beneath Norway Maple in Muttontown Preserve, Nassau County, which resulted in 80% of soil N being in the nitrate form, which can readily leach from the soil (unlike ammonium N). Fang's unpublished work needs to be verified in other locations. Fellows (2004) reports insignificant effects to ecosystem processes but the reference cited, Randall & Marinelli, 1996, does not address this issue in detail. Soil density may be increased, need sources or verification.

Sources of information:

Donnelly, CIPWG undated; Abbey, 2000; Andrews pers. obs.; Clemants pers. obs.; Love, 2003; Fellows, 2004; Fang 2005; Fang unpublished; Swearingen et al., 2002; Weber, 2003.

1.2. Impact on Natural Community Structure

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score 10

Documentation:

Identify type of impact or alteration:
 Oftentimes will increase the density in the existing (tree) layer. It can also tolerate poor soils and extreme temperatures and thus leads to the creation of a new (tree) layer where

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

one might not have developed (or at least not as fast) were succession left to native species. Norway maple also suppresses the understory and reduces or eliminates the shrub and herb layer, and saplings of native tree species.

Sources of information:

Webb & Kaunzinger, 1993; Wyckoff & Webb, 1996; Webb et al., 2000; Love, 2003; Bertin et al., 2005; Reinhart et al. 2005; Martin & Marks, 2006; Galbraith-Kent & Handel. 2008.

1.3. Impact on Natural Community Composition

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) 10
- U. Unknown

Score

10

Documentation:

Identify type of impact or alteration:

Reduces species richness and biodiversity, suppresses growth of saplings of native tree species, suppresses or eliminates native shrub and herb species, and may facilitate growth of non-native plants.

Sources of information:

Webb & Kaunzinger, 1993; Webb et al., 2000; Wyckoff & Webb 1996; Love, 2003; Reinhart et al. 2005; Fang 2005; Galbraith-Kent & Handel, 2008 .

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades.

Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- A. Negligible perceived impact 0
- B. Minor impact 3
- C. Moderate impact 7
- D. Severe impact on other species or species groups 10
- U. Unknown

Score

10

Documentation:

Identify type of impact or alteration:

Species serves as a preferential host of Asian long-horned beetle (*Anoplophora glabripennis*), and other pest insect species. Large stands of Norway maple support high populations of these insect pests, which then can lead to infestations on less preferred native tree species. Fang's unpublished work on altered nitrification suggests the possibility of impacts on soil microbial populations, but needs confirmation.

Sources of information:

Gilman & Watson, 2003; Love, 2003; Fellows 2004; Fang unpublished.

Total Possible

40

Section One Total

37

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction (provisional thresholds, more investigation needed)

- A. No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). 0
- B. Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) 1
- C. Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) 2
- D. Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.) 4
- U. Unknown

Score

4

Documentation:

Describe key reproductive characteristics (including seeds per plant):
Copious production of viable seeds.

Sources of information:

Webb & Kaunzinger, 1993; Webb et al., 2000; Fellows, 2004; Meiners, 2005; Reinhart et al., 2005;

2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score

2

Documentation:

Identify dispersal mechanisms:

Seeds mainly fall within short distance from parent trees (less than 100 meters), but there is potential for long distance dispersal (100+ m.) by wind or water .

Sources of information:

Love, 2003; Fellows, 2004; Bertin et al., 2005; Invasive Species Specialist Group, 2006; author's (Moore's) personal observations.

2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Score 3

Documentation:
 Identify dispersal mechanisms:
 Sold commercially.
 Sources of information:
 Love, 2003; Fellows, 2004; Bertin et al., 2005; Invasive Species Specialist Group, 2006.

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 3
- C. Possesses two or more characteristics that increase competitive advantage 6
- U. Unknown

Score 6

Documentation:
 Evidence of competitive ability:
 Ability to grow on infertile soils, fast growth, shade tolerant, perennial. Allelopathy has been suspected but research indicates it is absent or minimal (Rice, 2004)
 Sources of information:
 Love, 2003; Fellows, 2004; Bertin et al., 2005; Invasive Species Specialist Group, 2006; Meiners, 2005; Reinhart et al., 2005; Webb et al., 2000 & 2001; Webster et al. 2005; Wyckoff & Webb 1996; authors' personal observations.

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
- B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
- U. Unknown

Score 0

Documentation:
 Describe growth form:
 Can form monoculture stands but not thickets.
 Sources of information:
 Love, 2003; Fellows, 2004; authors' personal observations.

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
- B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
- C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
- U. Unknown (No studies have been completed)

Score 3

Documentation:
 Describe germination requirements:
 Germinates in a wide variety of conditions.
 Sources of information:
 Wyckoff & Webb, 1996; Webb et al., 2000 & 2001; Bertin et al., 2005; Meiners, 2005; Reinhart et al., 2005; author's (Moore's) personal observations.

2.7. Other species in the genus invasive in New York or elsewhere

- A. No 0
- B. Yes 3

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

U. Unknown

Score

3

Documentation:

Species:

Acer pseudoplatanus.

Total Possible

25

Section Two Total

21

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: “The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude”)

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score

2

Documentation:

Identify reason for selection, or evidence of weedy history:

Large stands always present in disturbed landscapes, usually with other invasive species present. May be a function of time, with age, occurrences in natural areas may become dense stands.

Sources of information:

Webb & Kaunzinger, 1993; Webb et al., 2000 & 2001; Fellows, 2004; Webster et al., 2005; Wyckoff & Webb, 1996; Reinhart et al., 2005; Meiners, 2006; author's (personal observations).

3.2. Number of habitats the species may invade

- A. Not known to invade any natural habitats given at A2.3 0
- B. Known to occur in two or more of the habitats given at A2.3, with at least one a natural habitat. 1
- C. Known to occur in three or more of the habitats given at A2.3, with at least two a natural habitat. 2
- D. Known to occur in four or more of the habitats given at A2.3, with at least three a natural habitat. 4
- E. Known to occur in more than four of the habitats given at A2.3, with at least four a natural habitat. 6
- U. Unknown

Score

2

Documentation:

Identify type of habitats where it occurs and degree/type of impacts:

See A2.3

Sources of information:

Fellows, 2004; Brooklyn Botanic Garden, 2008.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

3.3. Role of disturbance in establishment

- A. Requires anthropogenic disturbances to establish. 0
- B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
- C. Can establish independent of any known natural or anthropogenic disturbances. 4
- U. Unknown

Score 2

Documentation: Identify type of disturbance: Rarely will establish in areas with natural disturbances (e.g., windfalls) but usually occurs in areas with anthropogenic disturbance. Sources of information: Love, 2003; Fellows, 2004; Reinhart et al., 2005, Webb et al., 2000 & 2001; Wyckoff & Webb, 1996; author's (Moore's) personal observations.	
--	--

3.4. Climate in native range

- A. Native range does not include climates similar to New York 0
- B. Native range possibly includes climates similar to at least part of New York. 1
- C. Native range includes climates similar to those in New York 3
- U. Unknown

Score 3

Documentation: Describe what part of the native range is similar in climate to New York: Europe. Sources of information: Invasive Species Specialist Group 2006; Brooklyn Botanic Garden, 2008.	
--	--

3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope)

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score 4

Documentation: Identify states and provinces invaded: CT, DC, DE, IL, IN, KY, MA, MD, ME, MI, MN, NH, NJ, NY, OH, PA, RI, TN, VA, VT, WI, WV; NB, NF, NS, ON, PE, QC. Sources of information: <ul style="list-style-type: none"> • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces. U.S.D.A., 2008.	
---	--

3.6. Current introduced distribution of the species in natural areas in the eight New York State PRISMs (Partnerships for Regional Invasive Species Management)

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- A. Present in none of the PRISMs 0
- B. Present in 1 PRISM 1
- C. Present in 2 PRISMs 2
- D. Present in 3 PRISMs 3
- E. Present in more than 3 PRISMs or on the Federal noxious weed lists 4
- U. Unknown

Score

Documentation:
 Describe distribution:
 Present in all PRISMs.
 Sources of information:
 Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008.

Total Possible
 Section Three Total

4. DIFFICULTY OF CONTROL

4.1. Seed banks

- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
- B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
- C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
- U. Unknown

Score

Documentation:
 Identify longevity of seed bank:
 Seeds remain viable for over a year but no evidence for 10 years.
 Sources of information:
 Wyckoff & Webb, 1996; Webb et al., 2001; Bertin et al., 2005; author's (Moore's) personal observations

4.2. Vegetative regeneration

- A. No regrowth following removal of aboveground growth 0
- B. Regrowth from ground-level meristems 1
- C. Regrowth from extensive underground system 2
- D. Any plant part is a viable propagule 3
- U. Unknown

Score

Documentation:
 Describe vegetative response:
 Sprouts from base after cutting.
 Sources of information:
 Swearingen et al., 2003; author's personal observations.

4.3. Level of effort required

- A. Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. 0
- B. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft²). 2

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- C. Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). 3
- D. Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). 4
- U. Unknown

Score

4

Documentation:

Identify types of control methods and time-term required:

Large stands of trees require a major effort of person hours; follow up needed for removal of seedlings and saplings from seed bank.

Sources of information:

Abbey, 2000; Webb et al., 2000, 2001; Bertin et al., 2005; Meiners, 2005; Reinhart et al., 2005.

Total Possible

10

Section Four Total

7

Total for 4 sections Possible

100

Total for 4 sections

82

C. STATUS OF CULTIVARS AND HYBRIDS:

At the present time (May 2008) there is no protocol or criteria for assessing the invasiveness of cultivars independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain, and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Some cultivars of the species known to be available: ‘Almira’, ‘Cleveland’, ‘Columnare’, ‘Crimson King’, ‘Deborah’, ‘Drummondii’, ‘Emerald Queen’, ‘Erectum’, ‘Globosum’, ‘Greenlace’, ‘Jade Glen’, ‘Parkway’, ‘Olmstead’, ‘Summershade’, ‘Superform’, ‘Royal Red’.

References for species assessment:

Abbey, T. 2000. Invasive Plant Information Sheet – Norway Maple <http://www.hort.uconn.edu/cipwg/docs/norway_maple.pdf> [accessed February 26, 2008].

Bertin, R. I., M. E. Manner, B. F. Larrow, T. W. Cantwell, & E. M. Berstene 2005. Norway maple (*Acer platanoides*) and other non-native trees in urban woodlands of central Massachusetts. *Journal of Torrey Botanical Society* 132(2): 225-235.

Brooklyn Botanic Garden. 2008. AILANTHUS database. [Accessed on February 26, 2008].

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

Donnelly, C. Connecticut Invasive Plant Working Group. undated. Norway maples. *Acer platanoides*. <www.hort.uconn.edu/cipwg/invader_month/acerplat1.pdf>. [Accessed September 4, 2008.]

Fang, W. 2005. Spatial analysis of an invasion front of *Acer platanoides*: dynamic inferences from static data. *Ecography* 28(3): 283-294.

Fang, W. Unpublished. Is Norway maple a threat to Long Island's forests? PowerPoint presentation to LIISMA 2004.

Fellows, M. 2004. *Acer platanoides*. U.S. Invasive Species Impact Rank (I-Rank). NatureServe Explorer <www.natureserve.org>. [Accessed on September, 2, 2008.]

Galbraith-Kent, S.L. and S.N. Handel. 2008. Invasive *Acer platanoides* inhibits native sapling growth in forest understorey communities. *Journal of Ecology* 96(2): 293-302.

Gilman, E. F. and D. G. Watson. 1993. *Acer platanoides* Norway Maple. Fact Sheet ST-28, adapted from a series by the Environmental Horticulture Department, University of Florida for the United States Forest Service.

Hong, T. D. & R. H. Ellis, 1990. A comparison of maturation drying, germination, and desiccation tolerance between developing seeds of *Acer pseudoplatanus* L. and *Acer platanoides* L. *New Phytologist*, Vol. 116 (4): 589-596.

Invasive Species Specialist Group 2006. Global Invasive Species Database <<http://www.issg.org/database/species/search.asp?sts=sss&st=sss&fr=1&sn=acer+platanoides&rn=&hci=-1&ei=-1&x=37&y=5>>. [accessed February 26, 2008].

Love, R. 2003. Introduced Species Summary Project Norway Maple *Acer platanoides*. Invasion Biology Introduced Species Summary Project - Columbia University. <www.columbia.edu/itc/cerc/danoff-burg/invasion_bio/inv_spp_summ/Acer_platanoides_2.htm>. [Accessed September, 2, 2008.]

Martin, P. H. & P. L. Marks, 2006. Intact forests provide only weak resistance to a shade-tolerant invasive Norway maple (*Acer platanoides* L.). *Journal of Ecology* 94(6): 1070-1079.

Meiners, S. J. 2005. Seed and seedling ecology of *Acer saccharum* and *Acer platanoides*: A contrast between native and exotic congeners. *Northeastern Naturalist* 12(1): 23-32.

New York Flora Association. 2008. New York Flora Atlas. <<http://atlas.nyflora.org/>> [Accessed February 26, 2008].

Randall, J.M. and J. Marinelli (eds.) 1996. *Invasive Plants: Weeds of the Global Garden*. Brooklyn Botanic Garden, New York. 112 pp.

Rice, E. L. 2004. Investigation of allelopathy in an invasive introduced tree species, Norway maple (*Acer platanoides* L.). Doctoral thesis. DrexelUniversity. 148 pp. <http://dspace.library.drexel.edu/bitstream/1860/294/6/rich_elizabeth_thesis.pdf>. [Accessed September 4, 2008.]

Reinhart, K. O., E. Greene, and R. M. Callaway. 2005. Effects of *Acer platanoides* invasion on understory plant communities and tree regeneration in the northern Rocky Mountains. *Ecography* 28(5):573-582.

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

Swearingen, J., K. Reshetiloff, B. Slattery, and S. Zwicker. 2002. Plant Invaders of Mid-Atlantic Natural Areas. National Park Service and U.S. Fish & Wildlife Service, 82 pp.

United States Department of Agriculture, National Resources Conservation Service. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana [Accessed on February 26, 2008].

Wangen, S. R. & C. R. Webster, 2006. Potential for multiple lag phases during biotic invasions: reconstructing an invasion of the exotic tree *Acer platanoides*. *Journal of Applied Ecology* 43(2): 258-268.

Wangen, S. R., C. R. Webster, & J. A. Griggs, 2006. Spatial characteristics of the invasion of *Acer platanoides* on a temperate forested island. *Biological Invasions* 8(5): 1001-1012.

Webb, S. L. & C. K. Kaunzinger 1993. Biological invasion of the Drew University (New Jersey) Forest Preserve by Norway maple (*Acer platanoides* L.). *Bulletin of Torrey Botanical Club* 120(3): 343-349.

Webb, S. L., M. Dwyer, C. K. Kaunzinger & P. H. Wyckoff 2000. The myth of the resilient forest: Case study of the invasive Norway maple (*Acer platanoides*). *Rhodora* 102(911): 332-354.

Webb, S. L., T. H. Pendergast IV & M. F. Dwyer 2001. Response of native and exotic maple seedling banks to removal of the exotic, invasive Norway maple (*Acer platanoides*). *Journal of Torrey Botanical Society* 128(2): 141-149.

Weber, E. 2003. Invasive plant species of the world: a reference guide to environmental weeds. CABI Publishing, Cambridge, Massachusetts. 548 pp.

Webster, C. R., K. Nelson, S. R. Wangen, 2005. Stand dynamics of an insular population of an invasive tree, *Acer platanoides*. *Forest Ecology and Management* 208(1-3): 85-99.

Wyckoff, P. H. & S. L. Webb 1996. Understory influence of the invasive Norway maple (*Acer platanoides*). *Bulletin of Torrey Botanical Club* 123(3): 197-205.

Weldy, T. and D. Werier. 2005. New York Flora Atlas. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. [Accessed on February 26, 2008].

Citation: This NY ranking form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

Acknowledgments: The NY form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form. Original members of the LIISMA SRC included representatives of the Brooklyn Botanic Garden; The Nature Conservancy; New York Natural Heritage Program, New York Sea Grant; New York State Office of Parks, Recreation and Historic Preservation; National Park Service; Brookhaven National Laboratory; New York State Department of Environmental Conservation Region 1; Cornell Cooperative Extension of Suffolk/Nassau Counties; Long Island Nursery and Landscape Association; Long Island Farm Bureau; SUNY Farmingdale Ornamental Horticulture Department; Queens College Biology Department; Long Island Botanical Society; Long Island Weed Information Management System database

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

manager; Suffolk County Department of Parks, Recreation and Conservation; Nassau County Department of Parks, Recreation and Museums; Suffolk County Soil & Water Conservation District.

References for ranking form:

- Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.
- Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).
- Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>
- Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49
- Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.
- Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.